FIAN/TD-19/00
ITEP-TH-66/00
hep-th/0011222

Seiberg-Witten Toda Chains and N=1 SQCD [1]

A.Marshakov [2]

Theory Department, Lebedev Physics Institute, Moscow 117924, Russia
and
ITEP, Moscow 117259, Russia,

arXiv:hep-th/0011222v2 30 Nov 2000

We consider the Seiberg-Witten Toda chains arising in the context of exact solutions to $\mathcal{N}=2$ SUSY Yang-Mills and their relation to the properties of $\mathcal{N}=1$ SUSY gauge theories. In particular, we discuss their "perturbative" and "solitonic" degenerations and demonstrate some relations of the latter ones to the confining properties of $\mathcal{N}=1$ vacua.

1 Introduction

The $\mathcal{N}=2$ SUSY gauge theories possess continuous parametric families of vacua, labeled by VEV's of complex scalar fields Φ. An exciting result is that the masses of BPS states and low-energy effective actions for massless fields in these theories can be found exactly [1] and are described in terms of integrable systems so that the moduli of $\mathcal{N}=2$ theories are identified with moduli of complex structure of the corresponding spectral curves [1, 2] or integrals of motion of completely integrable models [3]. The corresponding family of integrable models in the most simple case consists of the Toda chains [4] of exponentially interacting particles.

These $\mathcal{N}=2$ SUSY gauge theories are even relatively realistic from the point of view of their physical spectra, containing propagating massless abelian gauge fields, but, as a consequence of extended supersymmetry they possess also massless scalars – the excitations of moduli, whose presence contradicts to what one should expect in reality. The situation is seriously improved when $\mathcal{N}=2$ SUSY is violated down to $\mathcal{N}=1$, then in contrast to $\mathcal{N}=2$ situation, in $\mathcal{N}=1$ case a superpotential $W(\Phi)$ may be generated which fixes the VEV of scalars to the extrema $dW=0$ and one ends up with some restricted set (usually *finite* number) of vacua where there is no more place for light moduli scalar excitations. It is widely believed that $\mathcal{N}=1$ gauge theories are already almost the theories close to real QCD, therefore investigation of the properties of these vacua is an important problem.

Below, we will try to present some arguments that the Seiberg-Witten integrable systems, arose originally and associated usually to the $\mathcal{N}=2$ situation, may also play some role in investigating properties of $\mathcal{N}=1$ gauge theories. We will start with generic properties of SUSY algebra in supersymmetric Yang-Mills theories and demonstrate that already at this level one may expect appearance of certain hamiltonian and/or integrable structures. Then we will pass to direct investigation of particular limits of the Seiberg-Witten integrable systems, since it is clear from the above reasoning that more realistic $\mathcal{N}=1$ picture appears only after some conditions on moduli are imposed. We will discuss, in particular, *degenerations* of the smooth Seiberg-Witten curves and demonstrate that, in fact, these degenerations can be separated into two different classes, the properties of integrable systems in these limits were studied in [5]. It turns out, that the structure of an integrable system implies a presence of a kind of "regularizing" multiplet of *fundamental* matter – to be identified with the set of Baker-Akhiezer functions of an integrable model, which can be explicitly constructed and their properties will be discussed below. In particular, we will see that the perturbative Baker-Akhiezer functions satisfy the condition, which may be thought of as a (holomorphic square root of) the D-term condition for the fundamental matter. We complete our discussion by few remarks about the "internal" properties of the Toda chain solutions.

[1] Based on the talks given at the *E.S.Fradkin Memorial Conference, Moscow, June 2000* and NATO workshop *Integrable Hierarchies and Modern Physics Theories, Chicago, July 2000* .

[2] e-mail address: mars@lpi.ru, mars@gate.itep.ru, marshakov@nbivms.nbi.dk




Figure 1: *Smooth Seiberg-Witten curve for the $\mathcal{N}=2$ $SU(N)$ pure gauge theory associated with periodic Toda chain.*

2 Supersymmetry and integrable structures

The basic understanding of integrable structure in $\mathcal{N}=2$ SUSY gauge theories appears already at the level of the (extended) SUSY algebra. Indeed $\mathcal{N}=2$ SUSY algebra in 4D is generated by the supercurrents $Q_A=\int d^3xJ^0_A$ $(A=1,2)$ and their complex conjugated. These currents differ from each other by exchanging the fermions $\{\psi_A\}$ and have general structure

$$J^0_A=(\psi_A F)\oplus(\psi_A\mathbf{D})\oplus\epsilon_{AB}(\bar{\psi}_B D\Phi)\oplus\epsilon_{AB}(\psi_B\mathbf{F}) \tag{1}$$

where $F=DA$ is gauge field strength, $\mathbf{D}^a(\Phi^\dagger,\Phi)=[\Phi^\dagger,\Phi]^a$ is the D-term and $\mathbf{F}^a(\Phi)=\frac{\partial W}{\partial\Phi^a}$ is the F-term [3]. From this form and canonical commutation relations it is clear that the anticommutator of Q_1 and Q_2 would contain the following contributions

$$\{Q_1,Q_2\}=\sum_{A=1,2}\{(\bar{\psi}_A D\Phi),(\psi_A F)\}+\dots \tag{2}$$

and these terms lead to the central charges [6]

$$Z_{SUSY}=\frac{1}{g^2_{YM}}\int D\Phi\wedge(F+*F) \tag{3}$$

In the case of point-like charged objects the 3-dimensional integral in the last formula can be rewritten using the Stokes theorem as an integral over sphere $\mathbf{S}^2$ at spatial infinity, giving rise to $Z_{SUSY}\sim a(n_e+\tau n_m)$, where $a\sim\langle\Phi\rangle$ is VEV of scalar and $\tau=\vartheta+\frac{i}{g^2_{YM}}$ is (complexified) coupling constant. However, if we originally consider instead of $\mathbf{R}^3$ a compactified space $\mathbf{R}^2\times\mathbf{S}^1$, the 2-dimensional integral is rather taken over $\mathbf{S}^1\times\mathbf{S}^1$ and can be presented as

$$Z_{SUSY}=\int_{\mathbf{S}^1\times\mathbf{S}^1}d^2x\epsilon^{ij}\partial_i\Phi\partial_j A=\int_{2-cycle}\delta\Phi\wedge\delta A \tag{4}$$

The last integral may be, in fact, identified with the integral of *holomorphic symplectic* form $\delta\Phi\wedge\delta A$ where the appearance of extra moduli parameters A is related to the Wilson-Polyakov loops of gauge fields, complexified by dual photons and becomes clear if one considers compactification of 4D theory down to 3D plus possible one compact dimension [7]. The loops of gauge field break the invariance in colour space and, in particular, this leads to replacement of the diagonal matrix of the VEV Φ by, in the simplest case of Toda chains "almost diagonal", Lax operator $\mathcal{L}$ of an integrable model.

Moreover, in the compactified case one may violate $\mathcal{N}=2$ 4D SUSY by the Scherk-Schwarz mechanism, which leads to generation of superpotentials, in spirit of [8]. These superpotentials exactly coincide with the Hamiltonians of the Seiberg-Witten integrable systems [8, 9]. It means, in particular, that integrable models are even more visible in *compactified* $\mathcal{N}=1$ theory ($\mathcal{N}=2$ in 3D terms), than in 4D $\mathcal{N}=2$ case.

SUSY breaking from $\mathcal{N}=2$ down to $\mathcal{N}=1$ (in 4D terms) is usually associated with *degeneration* [1, 10, 11] of smooth Seiberg-Witten curve for $\mathcal{N}=2$, see fig. 1. In fact not *any* degeneration of a curve corresponds to $\mathcal{N}=1$ vacuum. Roughly speaking there are two classes of degenerations, the "open" or *perturbative* and *solitonic*, if speaking in the language of integrable systems, the same may be thought of as asymptotically free and zero-charge from the point of view of corresponding quantum field theories. Only the latter class of (solitonic

[3] The superpotential terms are, in fact, absent in $\mathcal{N}=2$ theory, but they arise, in general, after breaking SUSY down to $\mathcal{N}=1$, which is important in the discussion below.




Figure 2: *A perturbative degeneration of the Seiberg-Witten curve from fig. 1 shrinks into its solitonic limit.*

or zero-charge) degenerations may play the role of set of vacua for $\mathcal{N}=1$ SUSY gauge theories. Qualitatively the difference is clearly seen in the brane picture for the Seiberg-Witten Toda chain curves [12, 13, 14, 15], the brane with *positive* tension would like to "shrink" to the zero-charge points, even if you start from different, perturbative degeneration (see fig. 2).

3 Integrable Toda chains from weak coupling

We start from *perturbative* degeneration, where $\mathcal{N}=2$ SUSY effective actions and integrable systems behind are completely determined by the 1-loop contributions. In the example of pure $\mathcal{N}=2$ gauge theory with $SU(N)$ gauge group [2], the scalar field Φ acquires nonzero VEV's $\Phi = \mathrm{diag}(a_1,\ldots,a_N)$ extremizing the potential $\mathrm{Tr}[\Phi,\Phi^\dagger]^2$. This (generically) breaks the $SU(N)$ gauge group due to the interaction term $\left([A_\mu,\Phi]_{ij}\right)^2 = \left(A_\mu^{ij}(a_i-a_j)\right)^2$ and the masses of W-bosons (and their superpartners) are classically given by $a_{ij}\equiv a_i - a_j$ or

$$a_{ij} = \oint_{C_{ij}} dS^{\mathrm{pert}} = \oint_{C_{ij}} \lambda d\log w = \oint_{C_{ij}} \lambda d\log P_N(\lambda) \tag{5}$$

for a particular "figure-of-eight" like contour C_{ij} around the roots $\lambda = a_i$ and $\lambda = a_j$ of the generating polynomial

$$w = P_N(\lambda) = \det_{N\times N}(\lambda - \Phi) = \prod_{i=1}^{N}(\lambda - a_i), \tag{6}$$

This means that the contour integrals in the complex λ-plane (5) for such contours around these singular points (the roots of the polynomial (6)) give one set of the BPS masses in the Seiberg-Witten theory. Another set of masses (monopoles) may be associated to *dual* contours starting and ending at the points a_i where dS^{pert} has singularities, taking into account that divergences are absorbed by a renormalization of the couplings. The integrals over the contours on the marked plane are in fact contour integrals on a *degenerate* Riemann surface depicted at the top of fig. 2: if one cuts it (vertically) into two pieces, connected by very narrow tubes, one gets exactly two copies of the desired λ-plane.

The $\mathcal{N}=2$ perturbative effective action (or prepotential) $\mathcal{F}$ and the set of effective couplings $-i\pi T_{jk} = \frac{\partial^2\mathcal{F}}{\partial a_j \partial a_k}$, are given by the 1-loop diagrams with the result [4]

$$T_{jk}^{\mathrm{pert}} = \frac{1}{2\pi i}\sum_{(\mathrm{massesM})_{jk}} \log\frac{(\mathrm{mass})^2}{\Lambda^2} = \frac{1}{2\pi i}\log\frac{a_{jk}^2}{\Lambda^2} \tag{7}$$

[4] Instanton contributions to the prepotential which are proportional to powers of Λ^{2N} are suppressed, and one keeps only the terms proportional to $\log\Lambda$ or the bare coupling τ.

where the scale parameter $\Lambda \equiv \Lambda_{QCD}$ so introduced may be related to the bare coupling τ.

The connection with integrable systems comes by interpreting the differential $dS^{\rm pert}$ with the generating differential of a Hamiltonian system – an open Toda chain or Toda molecule [5]. Remarkably, this interpretation extends beyond the perturbative regime, the "regularized" curve for (6) (see fig. 1!)

$$w + \frac{\Lambda^{2N}}{w} = P_N(\lambda) \tag{8}$$

is the spectral curve of a *periodic* Toda chain (an analog of the Sine-Gordon model in the same sense as the Toda molecule is a multiparticle analog of the Liouville model) providing the *exact* non-perturbative solution taking into account all instantonic contributions. The residue formulas (5) for the smooth curve (8) of genus $N-1$ turn into the action period integrals

$$(\mathbf{a}, \mathbf{a}^D) = \oint_{\mathbf{A},\mathbf{B}} \lambda \frac{dw}{w} \tag{9}$$

so that the set of effective charges (7) is generalized to the *period matrix* of (8)

$$T_{ij} = \frac{\partial a_i^D}{\partial a_j} = \frac{\partial^2 \mathcal{F}}{\partial a_i \partial a_j} \qquad i,j = 1,\ldots,N-1 \tag{10}$$

where $\mathcal{F}$ now stands for *non-perturbative* prepotential or effective action.

4 More on Toda: periodic chain and molecule

The dynamics of the integrable system itself may be again seen only in the compactified version [7] when the number of moduli of the gauge theory jumps up to the whole phase space of an integrable model. In this case the angle variables play the role of extra moduli related directly to the hamiltonian "times". This relation is especially simple in perturbative regime, when the basis of (normalized) "holomorphic" differentials

$$2\pi i d\omega_i = \frac{d\lambda}{\lambda - a_i} \tag{11}$$

and corresponding angle variables (originally defined as co-ordinates in the Jacobian of smooth curve (8) of fig. 1) are

$$\begin{aligned} 2\pi i z_i = \sum_{k=1}^{N} \int_{P_0}^{P_k} \frac{d\lambda}{\lambda - a_i} = \sum_{n=0}^{\infty} a_i^n \sum_{k=1}^{N} \int_{P_0}^{P_k} \frac{d\lambda}{\lambda^{n+1}} \\ \underset{\lambda_k \to \lambda(P_0)=\infty}{\sim} N\log\lambda + \sum_{n=1}^{\infty} a_i^n t_n \end{aligned} \tag{12}$$

for some divisor $\{P_k\}$ with $\lambda_k \equiv \lambda(P_k)$ and

$$t_n = -\frac{1}{n}\sum_{k=1}^{N} \lambda_k^{-n} \tag{13}$$

are the "hamiltonian" times in Miwa variables. It means, that after renormalization $z_i \to z_i - \frac{N}{2\pi i}\log\Lambda$ one gets

$$2\pi i z_i = a_i t + \sum_{n>1} a_i^n t_n \tag{14}$$

A natural ingredient of the integrable formulation is the set of the Baker-Akhiezer functions $\{\Psi_i\}$ (and $\{\tilde\Psi\}$), which can be thought of as introducing of "regularizing" fundamental matter. Like quark multiplets in SUSY gauge theories the Baker-Akhiezer functions always appear "in pairs" and carry a single index of lowest fundamental, $\mathbf{N}$ – for $\{\Psi_i\}$, or $\bar{\mathbf{N}}$ – for $\{\tilde\Psi_i\}$, representation of gauge group. Again, in 4D theory they are, up to

[5] From the simplest $SU(2)$ pure gauge theory [1], with the curve (6) $w = \lambda^2 - h$, $h = \frac{1}{2}{\rm Tr}\Phi^2 = a^2$ it becomes clear, that the masses (5) are defined by the contour integrals of $dS^{\rm pert} = \frac{\lambda d\lambda}{\lambda - \sqrt{h}} + \frac{\lambda d\lambda}{\lambda + \sqrt{h}}$, which can be interpreted as an *integration* of open Toda-Liouville model with Hamiltonian $h = p^2 + e^{2q}$, provided $\lambda \to p$, $w \to e^{2q}$. The same is true in general, that is, integration of the canonical differential $dS = 2pdq$ over the trajectories of the Toda chain solutions gives rise, for various (complexified) trajectories, to the BPS masses in the Seiberg-Witten theory [3].

gauge transformation, just the eigenvectors of diagonal matrix Φ, but in compactified version, when Φ should be replaced by the Lax operator $\mathcal{L}$, they become nontrivial.

On degenerate spectral curve (6) the set of the Baker-Akhiezer functions can be defined by the following analytic requirements [5]:

- This is the set of functions $\Psi_k = \Psi_k(\lambda)$, $k = 0, \ldots, N-1$ [6] which have exactly k zeroes on rational curve (6) and a single pole of order k at $\lambda = \infty$. For $k \geq N$ they can be defined by $\Psi_{k+N} = w\Psi_k$.
- These functions can be constructed as linear combinations of $\prod_{i_1<\ldots<i_k}(\lambda - a_{i_l}) = \prod_{j\in I}(\lambda - a_j)$ or, vice versa, certain linear combinations of Ψ_k vanish exactly at the chosen sets I of k points $a_{i_1}, \ldots, a_{i_k}$, $i_1 < \ldots < i_k$. One can see that the Baker-Akhiezer functions, indeed, satisfy for $e^{Z_i} \equiv e^{2\pi i z_i}\prod_{j\neq i}\frac{1}{a_{ij}}$

$$\sum_{i=1}^{N} e^{Z_i}\Psi_k(a_i) = (-)^{k+1}\sum_{|I|=k+1}\prod_{i\in I} e^{Z_i}\prod_{i,j\in I} a_{ij} = 0 \tag{15}$$

It is easy to see, that so defined Baker-Akhiezer functions are related to the tau-functions and can be written (up to normalization) as

$$\Psi_k(\lambda) = \lambda^k\ \Theta_k\left(z_i - \sum_n \frac{a_i^n}{n\lambda^n}\right) = \sum\nolimits^k \prod_{i\in I} e^{2\pi i z_i}(\lambda - a_i)\prod_{j\in\bar{I}}\frac{1}{a_{ij}} \tag{16}$$

where we write all a_{ij} in such order that $i < j$. The sum $\sum^k$ is taken over all sets of $|I| = k$ indices with $\bar{I}$ denoting the complementary set to I. The tau-functions

$$\begin{aligned}\Theta_k = \sum_{\sum_{i=1}^N n_i = k;\ \sum_{i<j}(n_i-n_j)^2 = k(N-k)} e^{2\pi i\sum_{j=1}^N n_j z_j - i\pi\sum_{i<j=1}^N (n_i-n_j)^2 T_{n_i n_j}} = \\ = \sum\nolimits^k \prod_{i\in I} e^{2\pi i z_i}\prod_{j\in\bar{I}}\frac{1}{a_{ij}}\end{aligned} \tag{17}$$

are nothing but degenerate theta-functions for the Toda molecule system where in the perturbative limit $\Lambda \to 0$ the period matrix T_{ij} is given by (7) with the terms proportional to $\log\Lambda$ absorbed into some renormalizations: $z_i \to z_i - \frac{N}{2\pi i}\log\Lambda$, $\Theta_k \to \Lambda^{k^2}\Theta_k$ [7]. They can be obtained as degenerations of the theta-functions on (Jacobian of) smooth curve (8) and their properties can be found, for example, in [5]. The formulas (16) can be written down even more explicitly, say for the $N = 2$ case one gets

$$\begin{aligned}\Psi_1(\lambda) = \frac{\lambda}{a}\left(\exp\left(z - \sum_n \frac{a^n}{n\lambda^n}\right) + \exp\left(-z - \sum_n (-)^n\frac{a^n}{n\lambda^n}\right)\right) = \\ = \frac{e^z + e^{-z}}{a}\left(\lambda - a\tanh z\right) = e^{-q}(\lambda - p)\end{aligned} \tag{18}$$

providing the relations $\Theta_1 = \frac{1}{a}(e^z + e^{-z}) = e^{-q}$ and $p = a\tanh z$ [8]. One may also easily check that the vector

$$\Psi(\lambda) = \begin{pmatrix} 1 \\ \Psi_1(\lambda)\end{pmatrix} \tag{19}$$

is an eigenvector of the open Toda Lax operator $\mathcal{L}\Psi = \lambda\Psi$

$$\mathcal{L} = \begin{pmatrix} p & e^q \\ e^q & -p\end{pmatrix} \tag{20}$$

provided by $\lambda^2 = p^2 + e^{2q} = h = a^2$, which is related to the VEV of 4D scalar Φ by $\mathcal{L} = U\Phi U^{-1}$, where matrix U is constructed from the Baker-Akhiezer functions.

[6] The N-dimensional vectors like $\{\Psi_k\}$ are labeled here for simplicity of notations as $k = 0, \ldots, N-1$, unlike usual conventions in group theory.

[7] The last renormalization could be understood as a logarithmic renormalization of the bare coupling τ.

[8] Note, that $\Psi_2 = \lambda^2 - a^2 = 0$ when $\lambda = \pm a$ belongs for a spectrum of the Toda molecule. This is a common feature for all $\Psi_N = \prod_{i=1}^N(\lambda - a_i)$.

The *dual* Baker-Akhiezer functions $\{\tilde{\Psi}_k(\lambda)\}$ (which may be thought as coming from the "dual" sheet of degenerate Toda chain curve [16]) can be written, up to normalization, again in terms of the theta-functions as

$$\tilde{\Psi}_{N-k-1}(\lambda) = \lambda^{-k}\ \Theta_k\left(z_i + \sum_n \frac{a_i^n}{n\lambda^n}\right) = \sum{}^{k} \prod_{i\in I} \frac{e^{2\pi i z_i}}{\lambda - a_i} \prod_{j\in \bar{I}} \frac{1}{a_{ij}} \tag{21}$$

For example

$$\begin{gathered} \tilde{\Psi}_{N-1} = \frac{1}{\prod_{i=1}^N (\lambda - a_i)} \\ \tilde{\Psi}_{N-2}(\lambda) = \lambda^{-1}\ \Theta_1\left(z_i + \sum_n \frac{a_i^n}{n\lambda^n}\right) = \sum_{i=1}^N \frac{e^{2\pi i z_i}}{\lambda - a_i} \prod{}' \frac{1}{a_{ij}} = \frac{\sum_{i=1}^N e^{2\pi i z_i} \prod'_{j\neq i} \frac{\lambda - a_i}{a_{ij}}}{\prod_{i=1}^N (\lambda - a_i)} \\ \ldots \end{gathered} \tag{22}$$

so that for the $N = 2$ case one gets

$$\begin{gathered} \tilde{\Psi}_1(\lambda) = \frac{1}{\lambda^2 - a^2} \\ \tilde{\Psi}_0(\lambda) = \lambda^{-1}\ \frac{1}{a}\left(e^{z + \sum_n \frac{a^n}{n\lambda^n}} + e^{-z + \sum_n (-)^n \frac{a^n}{n\lambda^n}}\right) = \frac{\frac{e^z}{\lambda - a} + \frac{e^{-z}}{\lambda + a}}{a} = \\ = \left(\frac{e^z + e^{-z}}{a}\right)\ \frac{\lambda + a\tanh z}{\lambda^2 - a^2} = e^{-q}\frac{\lambda + p}{\lambda^2 - a^2} \end{gathered} \tag{23}$$

It is easy to see, for (18) and (23) in particular, that one may indeed impose *gluing* conditions

$$\Psi_i(a) = C\tilde{\Psi}_i(a) \tag{24}$$

for some coefficient C. In our basis (21) for the dual Baker-Akhiezer functions this coefficient has to be nontrivial, since $\tilde{\Psi}_k(a)$ are naively singular exactly in the points of the spectrum, but adjusting C, or changing the normalization of (21) we see, that the gluing condition (24) indeed holds.

These are the gluing conditions at weak coupling – and they may be thought of as taking holomorphic square root from the D-term condition for a single flavour of "regularizing" fundamental matter.

5 The strong coupling limit

The quantum moduli space of the 4D pure $SU(N)$ $\mathcal{N} = 2$ SYM has N maximally singular points at which $N-1$ monopoles become simultaneously massless – the confining vacua of an $\mathcal{N} = 1$ theory [1, 2, 10, 15], where the dual variables $a_i^D = \frac{\partial \mathcal{F}}{\partial a_i}$ are appropriate to describe the prepotential. The $\mathcal{N} = 2$ spectral curve (8) at these points is described in terms of a Chebyshev polynomial (in this section we put $\Lambda = 1$)

$$P_N^{\text{Chebyshev}}(\lambda) = 2\cosh z. \tag{25}$$

with a solution $\lambda = 2\cosh\left(\frac{z}{N} + i\frac{2\pi k}{N}\right)$ $(k = 0, \ldots N-1)$. These are the $\mathcal{N} = 1$ points of the theory, related by a $\mathbf{Z}_N$ symmetry. The hyperelliptic form of the spectral curve (8) (recall $y = w - 1/w$) is now

$$y^2 = P_N(\lambda)^2 - 4 = (\lambda^2 - 4)\prod_{j=1}^{N-1}(\lambda - 2\cos\frac{\pi j}{N})^2 \tag{26}$$

which is a "solitonic" curve in the *periodic* Toda chain [18, 5]: now the corresponding $\mathbf{B}$ periods have collapsed, in contrast to the perturbative case. Introducing

$$\lambda = 2\cosh\frac{z}{N} \equiv \xi + \xi^{-1}. \tag{27}$$

one maps the 2-sheeted cover of the λ-plane $y = \sqrt{\lambda^2 - 4}$ to a cylinder with co-ordinate ξ. Thus eqs. (25), (26) describe analytically a cylinder with $N-1$ distinguished pairs of points – an inside-out S-duality transform of the bottom curve from fig. 2. It is worth noting that in contrast to perturbative regime (the upper curve on

fig. 2) now a curve does *not* separate into two pieces. The canonical basis of "holomorphic" differentials in this limit can be can be chosen as ($j = 1, \ldots, N-1$)

$$d\omega_j^D = \frac{\sin\frac{\pi j}{N}}{\pi}\frac{d\xi}{\left(\xi - e^{\frac{i\pi j}{N}}\right)\left(\xi - e^{-\frac{i\pi j}{N}}\right)} \quad \underset{\xi_j^{\pm}\equiv e^{\pm\frac{i\pi j}{N}}}{=} \quad \frac{1}{2\pi i} d\log\frac{\xi-\xi_j^+}{\xi-\xi_j^-}. \tag{28}$$

These differentials are normalized to the **B**-cycles, here the cycles around the marked points $\xi_j^{\pm}$,

$$\oint_{B_i} d\omega_j^D = \oint_{\xi_i^+} d\omega_j^D = -\oint_{\xi_i^-} d\omega_j^D = \delta_{ij}, \tag{29}$$

while certain of the **A**-periods ($\oint_{A_j} d\omega_j^D = \int_{\xi_j^-}^{\xi_j^+} d\omega_j^D$) diverge, the others ($j \neq k$) being given by [9]

$$T_{jk}^D = \oint_{A_j} d\omega_k^D = \frac{1}{2\pi i}\log\frac{\sin^2\frac{\pi}{2N}(j-k)}{\sin^2\frac{\pi}{2N}(j+k)} = \frac{1}{i\pi}\log\frac{\sin\frac{\pi}{2N}|j-k|}{\sin\frac{\pi}{2N}(j+k)}. \tag{30}$$

This is an exact formula for the set of effective couplings in strong-coupling regime, which has "typical solitonic" form. The appearance of solitons at strong coupling is a sort of "intuitive explanation" of the absence of the point with $\langle\Phi\rangle = 0$ since the solitonic points correspond, on one hand, to maximal degeneracy and, on the other hand, to minimum of superpotential, being an analog of the "BPS" first-order solutions in Seiberg-Witten Toda chains.

The angles at strong coupling are given by

$$z_j^D = \sum_{k=1}^{N-1}\int^{\xi_k} d\omega_j^D = \frac{1}{2\pi i}\sum_{k=1}^{N-1}\log\frac{\xi_k-\xi_j^+}{\xi_k-\xi_j^-} = \frac{1}{\pi}\sum_{n=1}^{\infty} t_n \sin\frac{\pi j n}{N} \tag{31}$$

where $t_n = \sum_k \frac{\xi_k^{\pm n}}{n}$ depending on whether $\{\xi_k\}$, $k = 1, \ldots, N-1$ go to 0 or to ∞. We observe also that the relation (31) coincides with the vacuum value of the string tension, or monopole condensate, proportional to the SUSY breaking parameters t_k [10, 15, 18, 19]. The detailed study of the properties of these strings are beyond the scope of this note, a discussion of related questions can be found in [20].

In contrast to the weak-coupling regime, the gluing conditions at the strong coupling are [18, 5]

$$\Psi_n(\xi_j^+) = \Psi_n(\xi_j^-) \qquad j = 1, \ldots, N-1. \tag{32}$$

which means that the Baker-Akhiezer function on a cylinder remembers that it originally came from a genus $N-1$ Riemann surface, and each pair of points $\xi_j^+, \xi_j^- = \frac{1}{\xi_j^+}$ corresponds to a degenerate handle. Note, that the form of condition (32) is different from (24) (and in fact, is even more similar to (15)) and relates the same Baker-Akhiezer functions in different points of the strong-coupling spectrum. There is no natural relation between Ψ and $\tilde{\Psi}$ in solitonic phase, which should be probably related to the properties of fundamental matter multiplets at strong coupling. We are going to return to this problem elsewhere.

6 More remarks on the Toda tau functions

In this section we will try to make few more remarks concerning the degenerate limits of the Toda chain, which do not have at the moment direct relation to the properties of $\mathcal{N} = 1$ SQCD, nevertheless being of some interest by themselves. We believe, however, that their physics origin is related to the bulk/boundary duality between SUSY gauge theories and gravity.

Commutativity of the theta-functions.

The theta-functions in (17) may be obtained as degenerate limit of the coefficients Θ_k in the expansion [21] of the genus N theta function

$$\Theta_N = \sum_{k\in\mathbb{Z}_N} e^{2\pi i\frac{k}{N}z}\Theta_k(\mathbf{z}|T) \tag{33}$$

[9] This formula was independently proposed in [17].

We may already deduce an interesting consequence: the ratios of the coefficients $\Theta_k \equiv \Theta_k(\mathbf{z}|T)$ Poisson commute (see [22] and references therein),

$$\left\{\frac{\Theta_i}{\Theta_j}, \frac{\Theta_{i'}}{\Theta_{j'}}\right\} = 0, \qquad \forall\ i, j, i', j'. \tag{34}$$

The Poisson bracket here is that corresponding to the symplectic form

$$\Omega^{Toda} = \sum_{i=1}^{N-1} dz_i \wedge da_i = \sum_{i=1}^{N-1} dq_i \wedge dp_i, \tag{35}$$

where q_i and p_i are co-ordinates and momenta of the Toda chain and we are working in the centre of mass frame. The Poisson commutativity of the ratios (34) follows from the solution of the periodic Toda chain [23, 24]

$$e^{q_i} = \frac{\Theta_i}{\Theta_{i-1}}, \qquad \frac{\Theta_i}{\Theta_j} = \prod_{k=j+1}^{i} e^{q_k} \tag{36}$$

(Here $\Theta_0 = \Theta_N$ and the Θ_k are the Toda chain tau-functions depending on the discrete time k, the number of the particle.) Now because the coordinates q_i obviously Poisson commute, $\{q_i, q_j\} = 0$, we deduce (34). This expression gives a precise formulation of the old expectation that the Toda chain tau-functions Poisson commute with each other. In the case of *periodic* Toda chain, however, only the *ratios* are commutative (theta-functions themselves, co-ordinates in projective space, are defined only up to common multiplication), but in the degenerate perturbative limit $\Theta_0 = \Theta_N = 1$ and, thus, tau-functions commute themselves. Moreover, in the perturbative limit formulas (17) and (36) give an explicit solution to the Toda molecule [4] in terms of the action-angle variables.

Relation to matrix models.

It is widely known now that the partition functions of the matrix models are tau functions of Toda chains [25]. Expression (17), being a finite sum, still is a sort of partition function, when the sum is taken over the discrete spectrum of the finite Toda molecule. It is easy to demonstrate, that, indeed this tau function can be considered as a "discretized" analog of the partition function of hermitian 1-matrix model.

For example, one can verify that

$$\Theta_k = \det_{k\times k} K_{n+m}\Big|_{n,m=1,\ldots,k} \tag{37}$$

where the "moment matrix" $K_{nm} = K_{n+m}$ is "an average" w.r.t. Θ_1, i.e.

$$K_n = \langle a^n\rangle_1 = \sum_{i=1}^{N} e^{Z_i} a_i^{n-1}, \qquad \Theta_1 = \sum e^{Z_i} \tag{38}$$

(cf. with [25, 26]). The simplest proof [5] is based on the Cauchy-Binet formula for the $k \times N$ ($k \leq N$) rectangular matrices A_{ni} and B_{im} ($i = 1, \ldots, N$, $n, m = 1, \ldots, k$)

$$\det_{k\times k}\left(\sum_{i=1}^{N} A_{ni}B_{im}\right)\Big|_{n,m=1,\ldots,k} = \sum_{i_1<\ldots<i_k} \det_{k\times k} A_{ni}\Big|_{i\in I;\ n=1,\ldots,k} \det_{k\times k} B_{im}\Big|_{i\in I;\ n=1,\ldots,k}. \tag{39}$$

so that

$$\Theta_k = \sum\nolimits_{|I|=k} \prod_{i\in I} e^{2\pi i z_i} \prod_{j\in \bar{I}} \frac{1}{a_{ij}} = \sum\nolimits_{|I|=k} \prod_{i\in I} e^{Z_i} \prod_{i\neq j\in I} a_{ij} = \sum\nolimits_{|I|=k} \prod_{i\in I} e^{Z_i} \prod_{i<j;\ i,j\in I} a_{ij}^2 \tag{40}$$

is indeed a "discretized" analogue of the tau-function of 1-matrix model [26] since for *any* coefficients C_i

$$\begin{array}{l} \sum_{|I|=k} \prod_{i\in I} C_i \prod_{i<j;\ i,j\in I} a_{ij}^2 \equiv \sum_{I:\ i_1<\ldots<i_k} C_{i_1}\ldots C_{i_k} \prod_{i_n<i_m} (a_{i_n} - a_{i_m})^2 \\ = \sum_{i_1<\ldots<i_k} C_{i_1}\ldots C_{i_k}\ \det_{k\times k} a_i^{n-1}\Big|_{i\in I;\ n=1,\ldots,k} \det_{k\times k} a_i^{m-1}\Big|_{i\in I;\ m=1,\ldots,k} \\ = \det_{k\times k}\left(\sum_{i=1}^{N} C_i a_i^{n+m-2}\right)\Big|_{n,m=1,\ldots,k}. \end{array} \tag{41}$$

and, substituting $C_i = e^{Z_i}$, one arrives at eqs. (37), (38)

$$\Theta_k = \sum_{I:\ i_1<\ldots<i_k} e^{Z_{i_1}+\ldots Z_{i_k}} \prod_{i_n<i_m} (a_{i_n} - a_{i_m})^2 = \det_{k\times k}\left(\sum_{i=1}^{N} e^{Z_i} a_i^{n+m-2}\right)\Big|_{n,m=1,\ldots,k} = \det_{k\times k} K_{n+m}. \tag{42}$$

The "non-discretized" (though still "discrete") 1-matrix model should be restored in the $N \to \infty$ limit of the Toda molecule, under condition that the spectrum $\{a_i\}$ fills in the whole real axis. Then instead of the Cauchy-Binet formula (rectangular matrices become infinite in one of the direction) one can effectively use the technique of the orthogonal polynomials [25, 26].

7 Conclusion

We have discussed just a brief list of questions related to the problems of $\mathcal{N} = 1$ vacua is SUSY gauge theories. However, already at this level we can see that there are remarkable coincidence between the properties of $\mathcal{N} = 1$ theories and Seiberg-Witten Toda chains "at strong coupling". They include the relations between the Baker-Akhiezer functions and the formulas for the phases of solitons, which look like being "in charge" for confining strings in SUSY gauge theories. It is necessary also to point out that perturbative integrable system is "stable" under small changes of moduli a_i, while in strong coupling regime all moduli are completely "confined" to certain values.

Acknowledgements

I am indebted to H.Braden, I.Krichever, A.Vainshtein and A.Yung for many illuminating discussions on different subjects considered in this note and to V.Zaykin and A.Sorin for encouragement to write this contribution. The work was partially supported by RFBR grant No. 99-02-16122, INTAS grant No. 99-0590 and CRDF grant No. RF-2102 (6531) and grant for the support of scientific schools No. 00-15-96566.

References

[1] N.Seiberg and E.Witten, Nucl.Phys., **B426** (1994) 19, hep-th/9407087; Nucl.Phys., **B431** (1994) 484, hep-th/9408099.

[2] A.Klemm, W.Lerche, S.Theisen and S.Yankielowicz, Phys. Lett. **344B** (1995) 169; hep-th/ 9411048; hep-th/ 9412158;
P.Argyres and A.Faraggi, Phys. Rev. Lett. **73** (1995) 3931, hep-th/ 9411057.

[3] A.Gorsky, I.Krichever, A.Marshakov, A.Mironov and A.Morozov, Phys.Lett., **B355** (1995) 466-474, hep-th/9505035.

[4] M.Toda, *Theory of Nonlinear Lattices*, Springer-Verlag, 1981.

[5] H.Braden and A.Marshakov, hep-th/0009060.

[6] E.Witten and D.Olive, Phys.Lett. **78B** (1978) 97.

[7] N.Seiberg and E.Witten, In *Saclay 1996, The mathematical beauty of physics*, 333-366, hepth/9607163.

[8] I.Affleck, J.Harvey and E.Witten, Nucl.Phys. **B206** (1982) 413.

[9] S.Katz and C.Vafa, Nucl.Phys., **B497** (1997) 196-204, hep-th/ 9611090;
O.Aharony, A.Hanany, K.Intriligator, N.Seiberg and M.Strassler, Nucl.Phys., **B499** (1997) 67-99, hep-th/ 9703110;
C.Vafa, Adv.Theor.Math.Phys. **2** (1998) 497-503, hep-th/ 9801139;
N. Dorey, JHEP 9907 (1999) 021, hep-th/9906011;
A.Marshakov, Prog.Theor.Phys.Suppl. 135 (1999) 1, hep-th/9906029.

[10] M.Douglas and S.Shenker, Nucl.Phys., **B447** (1995) 271-296, hep-th/9503163.

[11] A.Gorsky, A.Vainshtein and A.Yung, hep-th/0004087.

[12] E.Witten, Nucl.Phys. **B500** (1997) 3, hep-th/9703166.

[13] A.Marshakov, M.Martellini and A.Morozov, Phys.Lett. **B418** (1998) 294, hep-th/9706050;
A.Marshakov, hep-th/9709001.

[14] E.Witten, Nucl.Phys.**B507** (1997) 658, hep-th/9706109.

[15] A.Hanany, M.Strassler and A.Zafaroni, Nucl.Phys., **B513** (1998) 87-118, hep-th/9707244.

[16] I.Krichever and K.Vaninsky, hep-th/0010184.

[17] J.Edelstein, M.Gomez-Reino and M.Marino, hep-th/0006113.

[18] A.Marshakov, in *Continuum Advances in QCD 1998*, World Scientific, 1998;
see also
A.Marshakov, *Seiberg-Witten Theory and Integrable Systems*, World Scientific 1999.

[19] J.Edelstein, M.Marino and J.Mas, Nucl.Phys. B541 (1999) 671, hep-th/9805172.

[20] J.Edelstein, W.Fuertes, J.Mas and J.Guilarte, hep-th/0001184.

[21] H.W.Braden, A.Marshakov, A.Mironov and A.Morozov, Nucl. Phys. **B573** (2000) 553; hep-th/9906240.

[22] A.Marshakov, Phys. Lett. **B476** (2000) 420-426, hep-th/9912124.

[23] E.Date and S.Tanaka, Prog.Theor.Phys. **55** (1976) 457-465; Prog.Theor.Phys.Suppl. **59** (1976) 107-125;
see also [4].

[24] I.M.Krichever, Sov. Math. Surveys, **33** (1976) 215-216; see also Appendix by I.M.Krichever to
B.Dubrovin, Sov. Math. Surveys **36** (1981) N2, 12.

[25] A.Gerasimov, A.Marshakov, A.Mironov, A.Morozov and A.Orlov, Nucl.Phys.**B357** (1991) 565.

[26] S.Kharchev, A.Marshakov, A.Mironov, A.Orlov and A.Zabrodin, Nucl.Phys.**B366** (1991) 569-601.